APC Anti-Human CD40 Monoclonal Antibody

 Catalog Number
 Vial Size

 H10402-11G
 25 tests

 H10402-11H
 100 tests



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization


Clone	Isotype	Reactivity
G28.5	Mouse IgG1	Human

Description

CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection from apoptosis.

Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand). The G28.5 antibody has been reported to promote B cell proliferation in the presence of anti-IgM, IL-4 or PMA.

Illustration of Immunofluorescent Staining



Log Fluoresence Intensity

Human peripheral blood lymphocytes stained with APC anti-human CD40

Product Information

Conjugation: APC

Formulation: PBS pH 7.2, 0.09% NaN₃,

0.2% BSA

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used from 20 μL to 5 μL per 100 μL of peripheral blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Banchereau J, et al. 1994. Annu. Rev. Immunol. 12:881.
- [2] Foy T, et al. 1996. Annu. Rev. Immunol. 14:591.

For Research Use Only.