

## APC Anti-Mouse CD127 (IL-7R $\alpha$ ) Monoclonal Antibody



天津三箭生物技术股份有限公司  
Tianjin Sungene Biotech Co., Ltd.  
精准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
M11271-11A	25 $\mu$ g
M11271-11C	100 $\mu$ g

<b>Market</b>	400-621-0003 marketing@sungenebiotech.com
<b>Support</b>	022-66211636-8024 techsupport@sungenebiotech.com
<b>Web</b>	www.sungenebiotech.com

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.  
This product is guaranteed up to one year from purchase.


### Purified Antibody Characterization

Clone	Isotype	Reactivity
A7R34	Rat IgG2a	Mouse

### Description

CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor  $\alpha$  chain or IL-7R $\alpha$ . It forms a heterodimer with the common  $\gamma$  chain ( $\gamma$ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

### Illustration of Immunofluorescent Staining



C57BL/6 mouse splenocytes stained with PE anti-mouse CD3 and APC anti-mouse CD127

### Product Information

**Conjugation:** APC

**Formulation:** PBS pH 7.2, 0.09% NaN<sub>3</sub>, 0.2% BSA

**Concentration:** 0.2 mg/ml

**Storage:** Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

**Application:** Recommended Application: FC

**Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used  $\leq 0.25 \mu$ g /10<sup>6</sup> cells in 100  $\mu$ l). Since applications vary, the appropriate dilutions must be determined for individual use.

### References

- [1] Sudo T, et al. 1993. P. Natl. Acad. Sci. USA 90:9125.
- [2] Okuno Y, et al. 2001. P. Natl. Acad. Sci. USA 99:6246.
- [3] Pillai M, et al. 2004. Leukemia Lymphoma 45:2403.

**For Research Use Only.**