## **Biotin Anti-Human CD10 Monoclonal Antibody**



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Catalog NumberVial SizeH20103-08A25 ugH20103-08C100 ug

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

## **Purified Antibody Characterization**

Clone	Isotype	Reactivity
HI10a	Mouse IgG1	Human

## Description

CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins, bradykinin, angiotensins I and II, and other biologically active peptides.

**Product Information** 

Conjugation: Biotin

Formulation: PBS pH 7.2, 0.09% NaN<sub>3</sub>,

0.2% BSA

**Storage:** Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

**Application:** Recommended Application: FC

**Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used  $\leq 0.5$  µg per  $10^6$  cells in 100 µl volume or 100 µl of whole blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

## References

[1] Shipp M, et al. 1993. Blood 82:1052.

[2] Lu B, et al. 1995. J. Exp. Med. 181:2271.

For Research Use Only.