Biotin Anti-Mouse CD40 Monoclonal Antibody

 Catalog Number
 Vial Size

 M10401-08B
 50 μg

 M10401-08E
 500 μg



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
FGK4.5	Rat IgG2a	Mouse

Description

CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34+ hematopoietic progenitors. CD40 regulates B cell development/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface Iginduced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154(gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Product Information

Conjugation: Biotin

Formulation: PBS pH 7.2, 0.09% NaN₃,

0.2% BSA

Concentration: 0.5 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used ≤ 1.0 µg /10⁶ cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Barclay A, et al. 1997. The Leukocyte Antigen FactsBook Academic Press.
- [2] Bancherou J, et al. 1994. Annu. Rev. Immunol. 12:881.
- [3] Clark EA, et al. 1996. P. Natl. Acad. Sci. USA 83:4494.

For Research Use Only.