# **Biotin Anti-Mouse IL-17A Monoclonal Antibody**



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

 Catalog Number
 Vial Size

 M100I17-08B
 50 μg

 M100I17-08E
 500 μg

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

## **Purified Antibody Characterization**

Clone	Isotype	Reactivity
17F3	Mouse IgG1	Mouse

## Description

IL-17A is a cystine-linked homodimeric pro-inflammatory cytokine produced by  $TH_{17}$  cells,a distinct  $CD4^{\scriptscriptstyle +}$  T cell lineage. IL-17A stimulates the production of the pro-inflammatory cytokines IL-1 $\beta$ , TNF $\alpha$ , and IL-6. IL-17A also induces production of the neutrophil chemoattractants IL-8, CXCL1, and CXCL6 thereby bridging adaptive and innate immunity . IL-17A is intimately involved in mucosal immunity against bacterial infections and has a putative role in some autoimmune disorders . IL-17A effects appear to be exerted primarily through binding to the IL-17RA.IL-17A binding induces production of cytokines, chemokines and other proteins through activation of the ERK1/2 MAP kinase, PI3K/Akt, p38, and NF-κB pathways . Phosphorylation of some Jaks and Stats has been observed.

#### **Product Information**

Conjugation:Biotin

**Formulation:** PBS pH 7.2, 0.09% NaN<sub>3</sub>, 0.2%

**BSA** 

Concentration: 0.5mg/ml

**Storage:** Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.** 

Application: Recommended Application: FC

**Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used  $\leq 0.25$  µg /10<sup>6</sup> cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

### References

[1] Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anticytokine antibodies. J Immunol Methods. 1995;188(1):117-128.

For Research Use Only.