

FITC Anti-Human CD11c Monoclonal Antibody



天津三箭生物技术股份有限公司
Tianjin Sungene Biotech Co., Ltd.
精准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
H30111-02G	25 tests
H30111-02H	100 tests

Market	400-621-0003 marketing@sungenebiotech.com
Support	022-66211636-8024 techsupport@sungenebiotech.com
Web	www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.
This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
3.9	Mouse IgG1	Human*

*Cross-Reactivity: Chimpanzee, Baboon, African Green, Cynomolgus, Rhesus, Squirrel Monkey

Description

Clone 3.9 preferentially binds the activated form of CD11c, is specific for I domain of CD11c, and is able to partially block the binding of CD11c and ICAM-4.3.9 binding is divalent cation dependent. While analyzing blood, it is best to use heparin as the anti-coagulant and not EDTA. Since the ability of clone 3.9 to bind to its target is divalent cation dependent, the usage of EDTA as an anti-coagulant may be detrimental to staining due to its chelating properties.

Product Information

Conjugation: FITC

Formulation: Aqueous buffer, 0.09% NaN₃, may contain carrier protein/stabilize.

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used 20 µL to 5 µL /10⁶ cells or 100 µL of whole blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Petty H. 1996. Immunol. Today 17:209.
- [2] Springer T. 1994. Cell 76:301.
- [3] Ihanus E, et al. 2007. Blood 109:802-810.

For Research Use Only.