FITC Anti-Mouse LPAM-1 (Integrin α4β7) Monoclonal



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

 Catalog Number
 Vial Size

 M100L2-02B
 50 μg

 M100L2-02E
 500 μg

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
DATK32	Rat IgG2a	Mouse

Description

DATK32 antibody is specific for a combinatorial determinate of integrin $\alpha4\beta7complex.$ Integrin $\alpha4\beta7$ is composed of a 150 kD ($\alpha4$ or CD49d) and a 130 kD ($\beta7$) heterodimer, also known as CD49d/ $\beta7$ or LPAM-1. Belonging to the Ig superfamily, it is found on the majority of peripheral lymphocytes and subsets of thymocytes and bone marrow cells (including mast cell progenitors). Integrin $\alpha4\beta7$ binds its ligands, VCAM-1 (CD106), MAdCAM-1 and fibronectin, and plays an important role in lymphocytes adhesion and the direction of migration of blood lymphocytes to the intestine and associated lymphoid tissues.

Product Information

Conjugation: FITC

Formulation: PBS pH 7.2, 0.09% NaN₃,

0.2% BSA

Concentration: 0.5 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used $\leq 1.0~\mu g$ /10⁶ cells in 100 μl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Andrew DP, et al. 1994. J. Immunol. 153:3847.
- [2] Berlin C, et al. 1994. Cell 74:185.
- [3] Gurish MF, et al. 2001 J. Exp. Med. 194:1243.
- [4] Hamann A, et al. 1994. J. Immunol. 152:3282.

For Research Use Only.