LE/AF Purified Anti-Mouse CD86 Monoclonal Antibody

	天津三箭生物技术股份有限公司 Tianjin Sungene Biotech Co., Ltd.	
	精准 高效 穩定 Precision Efficient Stable	
Market	400-621-0003 marketing@sungenebiotech.com	
	marketing@sungenebiotech.com	
Support	022-66211636-8024	
	022-66211636-8024 techsupport@sungenebiotech.com	
Web	www.sungenebiotech.com	
1100	www.ourigenebioteen.com	

 Catalog Number
 Vial Size

 M10861-14B
 50 μg

 M10861-14E
 500 μg

 M10861-14F
 1 mg

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity	
GL1	Rat IgG2a	Mouse	

Description

CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is a ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce co-stimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Reported Applications

This GL1 antibody has been reported for use in blocking the mixed lymphocyte reaction in vitro and inhibit the priming of cytotoxic T lymphocytes in vivo (along with antibodies against B7-1). Additional reported applications (for the relevant formats) include: immunoprecipitation1, immunohistochemical staining of acetonefixed frozen sections, immunofluorescence microscopy, and in vivo and in vitro blocking of T cell responses. GL1 is not suitable for immunohistochemical staining of formalin-fixed paraffin sections.

Product Information

Production Method: Stirred tank fermentation

Medium: Hybridoma-SFM + 1%FCS + Gln + Gluc + P/S

Purification Method: Protein G

Concentration: 1 mg/ml

Endotoxin: < 2.00 EU/mg (LAL)

Purity: >95% (by SDS-PAGE)

Sterile: 0.2 µm Filtration

Formulated: PBS, pH7.2

Storage: Keep as concentrated solution.Store at 4°C as an undiluted liquid. For extended storage aliquot contents and freeze at-20°C or lower. Avoid cycles of freezing and thawing.

For Research Use Only.