PE Anti-Human CD11c Monoclonal Antibody

Catalog Number	Vial Size
H30111-09G	25 tests
H30111-09H	100 tests



Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization


Clone	Isotype	Reactivity	
3.9	Mouse IgG1	Human*	

*Cross-Reactivity: Chimpanzee, Baboon, African Green, Cynomolgus, Rhesus, Squirrel Monkey

Description

Clone 3.9 preferentially binds the activated form of CD11c, is specific for I domain of CD11c, and is able to partially block the binding of CD11c and ICAM-4. 3.9 binding is divalent cation dependent. While analyzing blood, it is best to use heparin as the anti-coagulant and not EDTA. Since the ability of clone 3.9 to bind to its target is divalent cation dependent, the usage of EDTA as an anti-coagulant may be detrimental to staining due to its chelating properties.

Illustration of Immunofluorescent Staining



Log Fluoresence Intensity Human peripheral blood granulocytes stained with PE anti-human CD11c

Product Information

Conjugation: PE

Formulation: Aqueous buffer, 0.09% NaN₃, may contain carrier protein/stabilize.

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used 20 μ L to 5 μ L /10⁶ cells or 100 μ L of whole blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

Schlossman S, et al.
Knapp W, et al.
McMichael A, et al. Eds.
Vainer B, et al. 2000.
Ottonello L, et al.
Asai A, et al. 2009.

For Research Use Only.