# PE Anti-Mouse CD152 Monoclonal Antibody

Catalog Number	Vial Size
M11521-09B	50 µg
M11521-09D	200 µg



**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.


#### **Purified Antibody Characterization**

Clone	Isotype	Reactivity	
UC10-4F10-11	Hamster IgG	Mouse	

#### Description

CD152, also known as CTLA-4 or Ly-56, is a 33 kD member of the immunoglobulin superfamily. It is expressed on activated T and B lymphocytes. CD152 is similar to CD28 in amino acid sequence, structure, and genomic organization and these two receptors share common B7 family counter-receptors (B7-1, B7-2). Whereas CD28 delivers a costimulatory signal in T cell activation, CTLA-4 negatively regulates cell-mediated immune responses. CD152 is thought to play a role in the induction and maintenance of immunological tolerance as well as the development of protective immunity and thymocyte regulation.

### Illustration of Immunofluorescent Staining



## **Product Information**

Conjugation: PE

**Formulation:** PBS pH 7.2, 0.09% NaN<sub>3</sub>, 0.2% BSA

Concentration: 0.2 mg/ml

**Storage:** Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.** 

Application: Recommended Application: FC

**Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used  $\leq 1.0 \ \mu g/10^6$  cells in 100  $\mu$ l). Since applications vary, the appropriate dilutions must be determined for individual use.

### References

- [1] Barclay A, et al. 1997. The Leukocyte Antigen FactsBook Academic Press.
- [2] Allison JP, et al. 1995. Science 270:932.
- [3] Waterhouse P, et al. 1995. Science 270:985.
- [4] Linsley PS, et al. 1991. J. Exp. Med. 174:561.

For Research Use Only.