

PE Anti-Mouse LPAM-1 (Integrin $\alpha 4\beta 7$) Monoclonal Antibody



天津三箭生物技术股份有限公司
Tianjin Sungene Biotech Co., Ltd.
精准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
M100L2-09B	50 μ g
M100L2-09D	200 μ g

Market | 400-621-0003
marketing@sungenebiotech.com

Support | 022-66211636-8024
techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.
This product is guaranteed up to one year from purchase.


Purified Antibody Characterization

Clone	Isotype	Reactivity
DATK32	Rat IgG2a	Mouse

Description

DATK32 antibody is specific for a combinatorial determinate of integrin $\alpha 4\beta 7$ complex. Integrin $\alpha 4\beta 7$ is composed of a 150 kD ($\alpha 4$ or CD49d) and a 130 kD ($\beta 7$) heterodimer, also known as CD49d/ $\beta 7$ or LPAM-1. Belonging to the Ig superfamily, it is found on the majority of peripheral lymphocytes and subsets of thymocytes and bone marrow cells (including mast cell progenitors). Integrin $\alpha 4\beta 7$ binds its ligands, VCAM-1 (CD106), MAdCAM-1 and fibronectin, and plays an important role in lymphocytes adhesion and the direction of migration of blood lymphocytes to the intestine and associated lymphoid tissues.

Illustration of Immunofluorescent Staining



C57BL/6 bone marrow lymphocytes were stained with APC anti-mouse B220 and PE anti-mouse LPAM-1

Product Information

Conjugation: PE

Formulation: PBS pH 7.2, 0.09% NaN_3 , 0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used $\leq 1.0 \mu\text{g} / 10^6$ cells in 100 μl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Andrew DP, et al. 1994. J. Immunol. 153:3847.
- [2] Berlin C, et al. 1994. Cell 74:185.
- [3] Gurish MF, et al. 2001 J. Exp. Med. 194:1243.
- [4] Hamann A, et al. 1994. J. Immunol. 152:3282.

For Research Use Only.