PE/Cy5 Anti-Human CD33 Monoclonal Antibody

Catalog Number Vial Size
H20331-35G 25 tests
H20331-35H 100 tests



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.


Purified Antibody Characterization

Clone	Isotype	Reactivity
HIM3-4	Mouse IgG1	Human

Description

CD33 is a 67 kD type I transmembrane glycoprotein also known as Siglec-3, gp67, and p67. It is a sialoadhesion immunoglobulin superfamily member expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes and hematopoietic stem cells. CD33 functions as a sialic acid-dependent cell adhesion molecule with carbohydrate/lectin binding activity.

Illustration of Immunofluorescent Staining



Log Fluoresence Intensity

Human peripheral blood cells stained with PE/Cy5 Mouse IgG1 Isotype Control (open histogram) or PE/Cy5 antihuman CD33 filled histogram).

Product Information

Conjugation: PE/Cy5

Formulation: PBS pH 7.2, 0.09% NaN₃,

0.2% BSA

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used from 20 μL to 5 μL per 100 μL of peripheral blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Favaloro E, et al. 1988. Br. J. Haematol. 69:163.
- [2] Freeman S, et al. 1995. Blood 85:2005.
- [3] Schlossman S, et al. Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.

For Research Use Only.