

PE/Cy7 Anti-Human HLA-DR Monoclonal Antibody



天津三箭生物技术股份有限公司
Tianjin Sungene Biotech Co., Ltd.
精准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
H200H7-17G	25 tests
H200H7-17H	100 tests

Market | 400-621-0003
marketing@sungenebiotech.com

Support | 022-66211636-8024
techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.
This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
HI159	Mouse IgG2a	Human

Description

HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4⁺ T cells.

Product Information

Conjugation: PE/Cy7

Formulation: PBS pH 7.2, 0.09% NaN₃, 0.2% BSA

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used from 20 μ L to 5 μ L per 100 μ L of peripheral blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Levacher M, et al. 1990. Clin. Exp. Immunol. 81:177.
- [2] Terstappen L, et al. 1990. J. Leukocyte Biol. 48:138.
- [3] Edwards JA, et al. 1986. J. Immunol. 137:490.
- [4] van Es A, et al. 1984. Transplantation 37:65.
- [5] O'Doherty U, et al. 1994. Immunology 82:487.
- [6] Thomas R, et al. 1994. J. Immunol. 153:4016.
- [7] Grouard G, et al. 1996. Nature 384:364.

For Research Use Only.