PE/Cy7 Anti-Mouse NK1.1 (CD161) Monoclonal Antibody



Market | 400-621-0003

marketing@sungenebiotech.com

Support | 022-66211636-8024

techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

 Catalog Number
 Vial Size

 M100N2-17A
 25 μg

 M100N2-17C
 100 μg

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Clone	Isotype	Reactivity
PK136	Mouse IgG2a	Mouse

Description

NK-1.1 surface antigen is encoded by the NKR-P1B/NKR-P1C gene, also known as CD161b/CD161c and Ly-55. It is expressed on NK cells and NK-T cells in some mouse strains, including C57BL/6, FVB/N, and NZB, but not AKR, BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL, and 129. Expression of NKR-P1C antigen has been correlated with lysis of tumor cells in vitro and rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN-γ production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.

Product Information

Conjugation: PE/Cy7

Formulation: PBS pH 7.2, 0.09% NaN₃,

0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged

exposure to light. Do not freeze.

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used ≤ 1.0 µg /10⁶ cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Carlyle, J.R., et al. 1999. J. Immunol. 162:5917.
- [2] Sentman, C.L., etal. 1989. Hybridoma 8:605.
- [3] Koo, G.C., et al. 1984. Hybridoma 3:301.
- [4] Sentman, C.L., et al. 1989. J. Immunol. 142:1847.
- [5] Koo, G.C., et al. 1986. J. Immunol. 137:3742.
- [6] Karlhofer, F.M., et al. 1991. J. Immunol. 146:3662.

For Research Use Only.