PE/Cy7 Anti-Mouse TCR γ/δ Monoclonal Antibody

| Catalog Number | Vial Size |
|----------------|-----------|
| M100T61-17A | 25 µg |
| M100T61-17C | 100 µg |



Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

| Clone | Isotype | Reactivity | |
|-------|-------------|------------|--|
| GL3 | Hamster IgG | Mouse | |

Description

T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR γ/δ belongs to the immunoglobulin superfamily, which is involved in the recognition of certain bacterial and tumor antigens bound to MHC class I. γ/δ TCR associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most γ/δ T cells are CD4⁻/CD8⁻ although some are CD8⁺. T cells expressing the γ/δ TCR have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease. It has been reported that γ/δ T cells also play a principal role in antigen presentation.

Product Information

Conjugation: PE/Cy7

Formulation: PBS pH 7.2, 0.09% NaN_3 , 0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used ≤ 0.25 µg /10⁶ cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- Skarstein K, et al. 1995. Immunology 81:497.
- [2] Harrison LC, et al. 1996. J. Exp. Med. 184:2167.
- [3] Wildner G, et al. 1996. Eur. J. Immunol. 26:2140.
- [4] Brandes M, et al. 2005. Science 309:264.

For Research Use Only.