PerCP Anti-Human HLA-DR Monoclonal Antibody

Vial Size
25 tests
100 tests



Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.


Purified Antibody Characterization

Clone	Isotype	Reactivity
HI159	Mouse IgG2b	Human

Description

HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4⁺ T cells.

Illustration of Immunofluorescent Staining



Log Fluorscence Intensity

Human peripheral blood lymphocytes stained with PerCP anti-humanHLA-DR

Product Information Conjugation: PerCP

Formulation: PBS pH 7.2, 0.09% NaN_{3,} 0.2% BSA

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used from 20 μ L to 5 μ L per 100 μ L of peripheral blood. Please check your vial). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Levacher M, et al. 1990. Clin. Exp. Immunol. 81:177.
- [2] Terstappen L, et al. 1990. J. Leukocyte Biol. 48:138.
- [3] Edwards JA, et al. 1986. J. Immunol. 137:490.
- [4] van Es A, et al. 1984. Transplantation 37:65.
- [5] O'Doherty U, et al. 1994. Immunology 82:487.
- [6] Thomas R, et al. 1994. J. Immunol. 153:4016.
- [7] Grouard G, et al. 1996. Nature 384:364.

For Research Use Only.