PerCP Anti-Mouse CD4 (YTS191) Monoclonal Antibody

Vial Size

25 µg

100 µg

	天津三箭生物技术股份有限公司 Tianjin Sungene Biotech Co., Ltd.
	精准 高效 稳定 Precision Efficient Stable
Market	400-621-0003 marketing@sungenebiotech.com
I	marketing@sungenebiotech.com
Support	022-66211636-8024
cappert	022-66211636-8024 techsupport@sungenebiotech.com
Web	www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Purified Antibody Characterization

Catalog Number

M10041-32A


M10041-32C

Clone	Isotype	Reactivity	
YTS191	Rat IgG2b	Mouse	

Description

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, Ick.

Illustration of Immunofluorescent Staining



FITC anti-mouse CD3

C57BL/6 mouse splenocytes stained with FITC anti-mouse CD3 and PerCP anti-mouse CD4

Product Information

Conjugation: PerCP

Formulation: PBS pH 7.2, 0.09% NaN₃, 0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution. Store at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis (The amount of the reagent is suggested to be used ≤ 0.25 µg /10⁶ cells in 100 µl). Since applications vary, the appropriate dilutions must be determined for individual use.

References

- [1] Barclay, A., et al. 1997. The Leukocyte Antigen FactsBook, Academic Press.
- [2] Bierer, B.E., et al. 1989. Annu. Rev. Immunol. 7:579.
- [3] Janeway, C.A. 1992. Annu. Rev. Immunol. 10:645.

For Research Use Only.