

PerCP Anti-Mouse TCR γ/δ Monoclonal Antibody

天津三箭生物技术股份有限公司
Tianjin Sungene Biotech Co., Ltd.
精准 高效 稳定 Precision Efficient Stable

Catalog Number	Vial Size
M100T61-32A	25 μ g
M100T61-32C	100 μ g

Market | 400-621-0003
marketing@sungenebiotech.com

Support | 022-66211636-8024
techsupport@sungenebiotech.com

Web | www.sungenebiotech.com

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.
This product is guaranteed up to one year from purchase.


Purified Antibody Characterization

Clone	Isotype	Reactivity
GL3	Hamster IgG	Mouse

Description

T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR γ/δ belongs to the immunoglobulin superfamily, which is involved in the recognition of certain bacterial and tumor antigens bound to MHC class I. γ/δ TCR associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most γ/δ T cells are CD4⁺/CD8⁻ although some are CD8⁺. T cells expressing the γ/δ TCR have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease. It has been reported that γ/δ T cells also play a principal role in antigen presentation.

Illustration of Immunofluorescent Staining



FITC anti-mouse CD3

C57BL/6 mouse splenocytes stained with
FITC anti-mouse CD3 and PerCP anti-mouse TCR γ/δ

Product Information

Conjugation: PerCP

Formulation: PBS pH 7.2, 0.09% NaN₃,
0.2% BSA

Concentration: 0.2 mg/ml

Storage: Keep as concentrated solution.
Store at 4°C and protected from prolonged
exposure to light. **Do not freeze.**

Application: Recommended Application: FC

Usage: Each lot of this antibody is quality
control tested by immunofluorescent staining
with flow cytometric analysis (The amount of
the reagent is suggested to be used ≤ 0.25
 μ g /10⁶ cells in 100 μ l). Since applications
vary, the appropriate dilutions must be
determined for individual use.

References

- [1] Skarstein K, et al. 1995. Immunology 81:497.
- [2] Harrison LC, et al. 1996. J. Exp. Med. 184:2167.
- [3] Wildner G, et al. 1996. Eur. J. Immunol. 26:2140.
- [4] Brandes M, et al. 2005. Science 309:264.

For Research Use Only.